

Functional Inequalities in Stratified Lie groups with Sobolev, Besov, Lorentz and Morrey spaces

Diego Chamorro*, Anca-Nicoleta Marcoci†, Liviu-Gabriel Marcoci‡

December 5, 2018

Abstract

When $p > 1$, using as base space classical Lorentz spaces associated to a weight from the Ariño-Muckenhoupt class B_p , we will study Gagliardo-Nirenberg inequalities. As a by-product we will also consider Morrey-Sobolev inequalities. These arguments can be generalized to many different frameworks, in particular the proofs are given in the setting of stratified Lie groups.

Keywords: Improved Sobolev inequalities; Sobolev spaces; Besov spaces; Classical Lorentz spaces; Stratified Lie groups.

Mathematics Subject Classification 46E35 ; 26D10 ; 46E30 ; 22E30

1 Introduction and presentation of the results

The aim of this article is to provide, in the setting of stratified Lie groups, a general proof for a particular type of improved Sobolev inequalities of the following general form

$$\|f\|_{\dot{W}^{s_1, q}} \leq C \|f\|_{\dot{W}^{s, p}}^\theta \|f\|_{\dot{B}_\infty^{-\beta, \infty}}^{1-\theta}, \quad (1)$$

where $f \in \dot{W}^{s, p} \cap \dot{B}_\infty^{-\beta, \infty}$. Here we write $\dot{W}^{s, p}$ for homogeneous Sobolev spaces and $\dot{B}_\infty^{-\beta, \infty}$ for homogeneous Besov spaces (see Section 3 below for precise definitions). The parameters s, s_1, p, q and β defining Sobolev and Besov spaces in the previous inequality are related by the conditions $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$, but they do not depend on the dimension and in this sense these inequalities are more general than classical Sobolev inequalities; of course the inequalities above are sharper than classical ones. Historically, the first proof in the Euclidean setting of these inequalities is due to P. Gérard, F. Oru and Y. Meyer [21] and is based on a Littlewood-Paley decomposition and interpolation results applied to dyadic blocks. Another proof of these inequalities using maximal function and Hedberg's inequality is given in [11].

*Laboratoire de Mathématiques et Modélisation d'Evry (LaMME) - UMR 8071. Université d'Evry Val d'Essonne, 23 Boulevard de France, 91037 Evry Cedex, France. email: diego.chamorro@univ-evry.fr

†Department of Mathematics and Computer Science. Technical University of Civil Engineering, Bucharest, Bld. Lacul Tei, no. 124, sector 2. Romania. email: anca.marcoci@utcb.ro

‡Department of Mathematics and Computer Science. Technical University of Civil Engineering, Bucharest, Bld. Lacul Tei, no. 124, sector 2. Romania. email: lmarcoci@instal.utcb.ro

Let us mention now that in the case of Lorentz spaces $L^{p,q}$, H. Bahouri and A. Cohen [4] proved the inequality

$$\|f\|_{L^{p,q}} \leq C \|f\|_{\dot{B}_q^{s,q}}^{q/p} \|f\|_{\dot{B}_q^{s-n/q,\infty}}^{1-q/p} \quad \text{with } \frac{1}{p} = \frac{1}{q} - \frac{s}{n}. \quad (2)$$

Remark that in this estimate, the index q defining the Lorentz and the Besov spaces is related to the parameters p , s and the dimension n . This inequality was generalized in the Euclidean setting by D. Chamorro & P-G. Lemarié-Rieusset [13] to other values of the parameter q using interpolation techniques and pointwise estimates. In a recent article, V.I. Kolyada and F.J. Pérez Lázaro [23] gave an interesting proof for inequalities of type (1) and (2) based on the use of rearrangement inequalities and the properties of the Gauss-Weierstrass kernel.

Motivated by the use of Lorentz spaces in these previous works, in our first theorem we will provide a generalization (in stratified Lie groups) of improved Sobolev inequalities of type (1) by considering weighted Lorentz-based Sobolev spaces defined as the set of measurable functions $f : \mathbb{G} \rightarrow \mathbb{R}$ such that the quantity $\|(-\Delta)^{\frac{s}{2}} f\|_{\Lambda^p(w)}$ is bounded where for $s > 0$ the fractional power of the Laplacian is defined in Section 2 below, and for $1 < p < +\infty$ the space $\Lambda^p(w)$ denotes the classical Lorentz space of functions introduced in [25] and [26] defined as

$$\Lambda^p(w) = \left\{ f : \|f\|_{\Lambda^p(w)} = \left(\int_0^{+\infty} f^*(t)^p w(t) dt \right)^{1/p} < +\infty \right\},$$

where w is a weight in \mathbb{R}_+ and f^* denotes the non-increasing rearrangement of f (see [5] for standard notations). Many of the properties of these spaces depend on the weight w : in particular, if $w = 1$ we have $\Lambda^p(w) = L^p$ and if $w(t) = t^{p/q-1}$, with $1 \leq q \leq +\infty$, we obtain $\Lambda^p(w) = L^{q,p}$, where $L^{q,p}$ are the usual Lorentz spaces. In this work we will consider the weighted Lorentz space $\Lambda^p(w)$ such that the weight w satisfies the B_p condition, the reason for this is given by the fact that M. A. Ariño and B. Muckenhoupt showed in [3] that this B_p condition characterizes the boundedness of the Hardy-Littlewood maximal operator on $\Lambda^p(w)$ and this particular property will be intensively used in our proofs. See Section 3 for definitions and [3], [8], [10] and [33] for more details and properties concerning these functional spaces.

In this direction of generalization, standard Sobolev inequalities have been studied by A. Cianchi [15] in the context of Orlicz-Sobolev spaces, but improved inequalities of the general type (3) presented in Theorem 1 below are, to the best of our knowledge, new.

Theorem 1 *Let \mathbb{G} a stratified Lie group. Let $s > 0$, $w \in B_p$ be a weight and let $f : \mathbb{G} \rightarrow \mathbb{R}$ be a function such that $(-\Delta)^{\frac{s}{2}} f \in \Lambda^p(w)(\mathbb{G})$ and $f \in \dot{B}_\infty^{-\beta,\infty}(\mathbb{G})$. Then we have the following version of improved Sobolev inequalities:*

$$\|(-\Delta)^{\frac{s_1}{2}} f\|_{\Lambda^q(w)} \leq C \|(-\Delta)^{\frac{s}{2}} f\|_{\Lambda^p(w)}^\theta \|f\|_{\dot{B}_\infty^{-\beta,\infty}}^{1-\theta}, \quad (3)$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

The choice of the weights in the B_p class is given by two important facts. First, these weights allow us to consider general functional spaces, in particular we can easily recover standard Lorentz spaces. Second, these weights ensure that maximal function is bounded in the spaces $\Lambda^p(w)$ for $1 < p < +\infty$, and this feature is crucial as the proof of Theorem 1 requires this property. Note in particular that

inequality (3) is different from inequality (2) since Lorentz-Sobolev spaces are not included in the scale of Besov spaces.

Since our proof of Theorem 1 relies essentially on a pointwise inequality and on the boundedness of the Hardy-Littlewood maximal operator, it is possible to give a related result replacing classical Lorentz spaces by Morrey spaces $\mathcal{M}^{p,a}$ which are a useful generalization of Lebesgue spaces. Classical Hardy-Littlewood-Sobolev inequalities were studied in this functional framework by D. Adams [1] and by F. Chiarenza & M. Frasca [14] and our next theorem is an improvement of these inequalities.

Theorem 2 *Let \mathbb{G} be a stratified Lie group. Let $s > 0$, $1 < p < +\infty$ and $0 \leq a < n$ and let f be a function such that $(-\Delta)^{\frac{s}{2}} f \in \mathcal{M}^{p,a}(\mathbb{G})$ and $f \in \dot{B}_{\infty}^{-\beta,\infty}(\mathbb{G})$. Then we have*

$$\|(-\Delta)^{\frac{s_1}{2}} f\|_{\mathcal{M}^{q,a}} \leq C \|(-\Delta)^{\frac{s}{2}} f\|_{\mathcal{M}^{p,a}}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta}, \quad (4)$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

The plan of this article is the following. In Section 2 we present our general framework which is given by stratified Lie groups. These groups are quite natural generalization of \mathbb{R}^n but they present some particularities that should be taken into account in the computations. In Section 3 we give the precise definition of all the functional spaces used in the previous inequalities and in Section ?? - 4 we present the proofs of Theorems ?? and Theorem 1 respectively. Finally, in Section 5 we give the proof of Theorem 2 and some variations of the previous results.

2 Stratified Lie groups: notation and basic properties

As said in the introduction, stratified Lie groups are natural generalizations of \mathbb{R}^n when considering general dilation structures. Although stratified Lie groups share common features with \mathbb{R}^n , there are some special points that must be taken into account: for example these groups are no longer abelian and this fact requires to be careful in some computations, furthermore from the geometric point of view, the inner geometric structure of these groups can be very different from the euclidean setting. It is then necessary to recall some basic facts about stratified Lie groups, for further information see [17], [18], [38], [35] and the references given there in.

We start with the notion of *homogeneous group* \mathbb{G} which is the data of \mathbb{R}^n equipped with a structure of Lie group and we will always suppose that the origin is the identity. We define a *dilation structure* by fixing integers $(a_i)_{1 \leq i \leq n}$ such that $1 = a_1 \leq \dots \leq a_n$ and by writing:

$$\begin{aligned} \delta_{\alpha} : \mathbb{R}^n &\longrightarrow \mathbb{R}^n \\ x &\longmapsto \delta_{\alpha}[x] = (\alpha^{a_1} x_1, \dots, \alpha^{a_n} x_n). \end{aligned} \quad (5)$$

We will often note αx instead of $\delta_{\alpha}[x]$ and α will always indicate a strictly positive real number.

Of course, the Euclidean space \mathbb{R}^n with its group structure and provided with its usual dilations (i.e. $a_i = 1$, for $i = 1, \dots, n$) is a homogeneous group. Here is another example: if $x = (x_1, x_2, x_3)$ is an element of \mathbb{R}^3 , we can fix a dilation by writing $\delta_{\alpha}[x] = (\alpha x_1, \alpha x_2, \alpha^2 x_3)$ for $\alpha > 0$. Then, the well suited group law with respect to this dilation is given by $x \cdot y = (x_1, x_2, x_3) \cdot (y_1, y_2, y_3) = (x_1 + y_1, x_2 + y_2, x_3 + y_3 + \frac{1}{2}(x_1 y_2 - y_1 x_2))$. Remark in particular that this group law is no longer

abelian. The triplet $(\mathbb{R}^3, \cdot, \delta)$ corresponds to the Heisenberg group \mathbb{H}^1 which is the first non-trivial example of a homogeneous group. The *homogeneous dimension* with respect to dilation structure (5) is given by $N = \sum_{1 \leq i \leq n} a_i$. We observe that it is always larger than the topological dimension n since

each integer a_i verifies $a_i \geq 1$ for all $i = 1, \dots, n$. For instance, in the Heisenberg group \mathbb{H}^1 we have $N = 4$ and $n = 3$ while in the Euclidean case these two concepts coincide. Now we will say that a function on $\mathbb{G} \setminus \{0\}$ is *homogeneous* of degree $\lambda \in \mathbb{R}$ if $f(\delta_\alpha[x]) = \alpha^\lambda f(x)$ for all $\alpha > 0$. In the same way, we will say that a differential operator D is homogeneous of degree λ if $D(f(\delta_\alpha[x])) = \alpha^\lambda (Df)(\delta_\alpha[x])$, for all f in operator's domain. In particular, if f is homogeneous of degree λ and if D is a differential operator of degree μ , then Df is homogeneous of degree $\lambda - \mu$. The presence of a dilation structure is one of most important features of stratified Lie groups and the homogeneity with respect to these dilations will play a useful role in our computations.

From the point of view of measure theory, homogeneous groups behave in a traditional way since Lebesgue measure dx is bi-invariant and coincides with the Haar measure, thus for any subset E of \mathbb{G} we will note its measure as $|E|$. This fact also allows us to define Lebesgue spaces in a classical way (see also Section 3 below). The convolution will be a very useful tool in our computations, and for two functions f and g on \mathbb{G} it is defined by

$$f * g(x) = \int_{\mathbb{G}} f(y)g(y^{-1} \cdot x)dy = \int_{\mathbb{G}} f(x \cdot y^{-1})g(y)dy, \quad x \in \mathbb{G}.$$

However, since the group law of a stratified Lie group is not necessarily commutative, we do not have in general the identity $f * g = g * f$ and we need to take care of this fact. Nevertheless, we have at our disposal Young's inequalities:

Lemma 2.1 *If $1 \leq p, q, r \leq +\infty$ such that $1 + \frac{1}{q} = \frac{1}{p} + \frac{1}{r}$. If $f \in L^p(\mathbb{G})$ and $g \in L^r(\mathbb{G})$, then $f * g \in L^q(\mathbb{G})$ and we have the inequality $\|f * g\|_{L^q} \leq \|f\|_{L^p} \|g\|_{L^r}$.*

A proof is given in [18]. A weak version of Young's inequalities will be stated in Section 3.

For a homogeneous group $\mathbb{G} = (\mathbb{R}^n, \cdot, \delta)$ we consider now its Lie algebra \mathfrak{g} whose elements can be conceived in two different ways: as *left*-invariant vector fields or as *right*-invariant vector fields. The left-invariant vectors fields $(X_j)_{1 \leq j \leq n}$ are determined by the formula

$$(X_j f)(x) = \left. \frac{\partial f(x \cdot y)}{\partial y_j} \right|_{y=0} = \frac{\partial f}{\partial x_j} + \sum_{j < k} q_j^k(x) \frac{\partial f}{\partial x_k},$$

where $q_j^k(x)$ is a homogeneous polynomial of degree $a_k - a_j$ and f is a smooth function on \mathbb{G} . By this formula one deduces easily that these vectors fields are homogeneous of degree a_j and we have $X_j(f(\alpha x)) = \alpha^{a_j}(X_j f)(\alpha x)$. We will note $(Y_j)_{1 \leq j \leq n}$ the right invariant vector fields defined in a totally similar way:

$$(Y_j f)(x) = \left. \frac{\partial f(y \cdot x)}{\partial y_j} \right|_{y=0}.$$

A homogeneous group \mathbb{G} is *stratified* if its Lie algebra \mathfrak{g} breaks up into a sum of linear subspaces $\mathfrak{g} = \bigoplus_{1 \leq j \leq k} E_j$ such that E_1 generates the algebra \mathfrak{g} and $[E_1, E_j] = E_{j+1}$ for $1 \leq j < k$ and $[E_1, E_k] = \{0\}$ and $E_k \neq \{0\}$, but $E_j = \{0\}$ if $j > k$. Here $[E_1, E_j]$ indicates the subspace of

\mathfrak{g} generated by the elements $[U, V] = UV - VU$ with $U \in E_1$ and $V \in E_j$. The integer k is called the *degree* of stratification of \mathfrak{g} . For example, on Heisenberg group \mathbb{H}^1 , we have $k = 2$ while in the Euclidean case $k = 1$.

We will suppose from now on that \mathbb{G} is **stratified** with homogeneous dimension¹ $N \geq 4$. Within this framework, we will fix once and for all the family of vectors fields

$$\mathbf{X} = \{X_1, \dots, X_m\},$$

such that $a_1 = a_2 = \dots = a_m = 1$ ($m < n$), then the family \mathbf{X} is a base of E_1 and generates the Lie algebra of \mathfrak{g} , which is precisely the Hörmander's condition (see [18] and [38]) and this particular choice ensures several important properties, in particular to the family \mathbf{X} is associated the Carnot-Carathéodory distance d which is left-invariant and compatible with the topology on \mathbb{G} (see [38] for more details) and for any $x \in \mathbb{G}$ we will denote by $|x| = d(x, e)$ and for $r > 0$ we form open balls by writing $B(x, r) = \{y \in \mathbb{G} : d(x, y) < r\}$. By simple homogeneity arguments we obtain that stratified Lie groups have polynomial volume growth since we have $|B(\cdot, r)| = r^N |B(\cdot, 1)|$.

The main tools of this paper depend on the properties of the gradient, the Laplacian and the associated heat kernel, but before introducing them, we make here some remarks on general vectors fields X_j and Y_j . Let us fix some notation: for any multi-index $I = (i_1, \dots, i_n) \in \mathbb{N}^n$, one defines X^I by $X^I = X_1^{i_1} \dots X_n^{i_n}$ and Y^I by $Y^I = Y_1^{i_1} \dots Y_n^{i_n}$, furthermore we denote by $|I| = i_1 + \dots + i_n$ the order of the derivation of the operators X^I or Y^I and $d(I) = a_1 i_1 + \dots + a_n i_n$ the homogeneous degree of these ones. Now, for $\varphi, \psi \in \mathcal{C}_0^\infty(\mathbb{G})$ we have the equality

$$\int_{\mathbb{G}} \varphi(x) (X^I \psi)(x) dx = (-1)^{|I|} \int_{\mathbb{G}} (X^I \varphi)(x) \psi(x) dx.$$

The interaction of operators X^I and Y^I with convolutions is clarified by the following identities:

$$X^I(f * g) = f * (X^I g), \quad Y^I(f * g) = (Y^I f) * g, \quad (X^I f) * g = f * (Y^I g). \quad (6)$$

Finally, one will say that a function $f \in \mathcal{C}^\infty(\mathbb{G})$ belongs to the Schwartz class $\mathcal{S}(\mathbb{G})$ if the following semi-norms are bounded for all $k \in \mathbb{N}$ and any multi-index I : $N_{k,I}(f) = \sup_{x \in \mathbb{G}} (1 + |x|)^k |X^I f(x)|$.

Remark 2.1 To characterize the Schwartz class $\mathcal{S}(\mathbb{G})$ we can replace vector fields X^I in the semi-norms $N_{k,I}$ above by right-invariant vector fields Y^I .

For a proof of these facts and for further details see [18] and [19].

We define now the *gradient* on \mathbb{G} from vectors fields of homogeneity degree equal to one (*i.e.* those composing the family \mathbf{X}) by fixing $\nabla = (X_1, \dots, X_m)$. This operator is of course left invariant and homogeneous of degree 1. The length of the gradient is given by the formula $|\nabla f| = ((X_1 f)^2 + \dots + (X_m f)^2)^{1/2}$. We also define the right invariant gradient $\tilde{\nabla} = (Y_1, \dots, Y_m)$, and using (6) we have the identity $(\nabla f) * g = f * (\tilde{\nabla} g)$. We define now the Laplacian we are going to work with. Let us notice that in this setting there is not a single way to build a Laplacian, see for example

¹The lower bound $N \geq 4$ corresponds to the homogeneous dimension of the Heisenberg group \mathbb{H}^1 , which is the simplest non-trivial stratified Lie group.

[19]. In this article we will use the Laplacian, denoted by \mathcal{J} , which is given from the family \mathbf{X} in the following way

$$\mathcal{J} = \nabla^* \nabla = - \sum_{j=1}^m X_j^2. \quad (7)$$

This is a positive self-adjoint, hypo-elliptic operator (since the family \mathbf{X} satisfies the Hörmander's condition), having as domain of definition $L^2(\mathbb{G})$. Its associated *heat operator* on $\mathbb{G} \times]0, +\infty[$ is given by $\partial_t + \mathcal{J}$. We recall now some well-known properties of the heat operator and its associated kernel.

Theorem 3 *There exists a unique family of continuous linear operators $(H_t)_{t>0}$ defined on $L^1 + L^\infty(\mathbb{G})$ with the semi-group property $H_{t+s} = H_t H_s$ for all $t, s > 0$ and $H_0 = Id$, such that:*

- 1) *the Laplacian \mathcal{J} is the infinitesimal generator of the semi-group $H_t = e^{-t\mathcal{J}}$;*
- 2) *H_t is a contraction operator on $L^p(\mathbb{G})$ for $1 \leq p \leq +\infty$ and for $t > 0$;*
- 3) *the semi-group H_t admits a convolution kernel $H_t f = f * h_t$ where $h_t(x) = h(x, t) \in \mathcal{C}^\infty(\mathbb{G} \times]0, +\infty[)$ is the heat kernel which satisfies the following points:*
 - (a) *$(\partial_t + \mathcal{J})h_t = 0$ on $\mathbb{G} \times]0, +\infty[$, and $h(x, t) = h(x^{-1}, t)$, $h(x, t) \geq 0$ and $\int_{\mathbb{G}} h(x, t) dx = 1$,*
 - (b) *h_t has the semi-group property: $h_t * h_s = h_{t+s}$ for $t, s > 0$ and we have $h(\delta_\alpha[x], \alpha^2 t) = \alpha^{-N} h(x, t)$,*
 - (c) *For every $t > 0$, $x \mapsto h(x, t)$ belong to the Schwartz class in \mathbb{G} .*
- 4) *For $f \in \mathcal{C}^\infty(\mathbb{G})$ and for $t > 0$ we have $\mathcal{J}H_t(f) = H_t \mathcal{J}(f)$.*

For a detailed proof of these and other important facts concerning the heat semi-group see [18] and [31].

To close this section we recall the definition of the Laplacian's fractional powers. If $s > 0$ we write

$$\mathcal{J}^s f(x) = \frac{1}{\Gamma(k-s)} \int_0^{+\infty} t^{k-s-1} \mathcal{J}^k H_t f(x) dt, \quad (8)$$

for all $f \in \mathcal{C}^\infty(\mathbb{G})$ with k an integer greater than s . The interaction between this operator and the heat kernel is given by the following lemma.

Lemma 2.2 *If $1 \leq p \leq +\infty$, for $s > 0$ and for $t > 0$ we have the estimate $\|\mathcal{J}^{\frac{s}{2}} h_t\|_{L^p} \leq Ct^{-\frac{s+N(1-\frac{1}{p})}{2}}$.*

See [31] for a proof.

3 Functional spaces

We give in this section the precise definition of all the functional spaces involved in Theorems ??, 1 and 2. In a general way, given a norm $\|\cdot\|_X$, we will define the corresponding functional space $X(\mathbb{G})$ by $\{f \in \mathcal{S}'(\mathbb{G}) : \|f\|_X < +\infty\}$. The constant that appear in this paper such as C may change from one occurrence to the next.

- **Lebesgue spaces** $L^p(\mathbb{G})$. For a measurable function $f : \mathbb{G} \rightarrow \mathbb{R}$ and for $1 \leq p < +\infty$ we define Lebesgue space by the norm $\|f\|_{L^p} = \left(\int_{\mathbb{G}} |f(x)|^p dx \right)^{1/p}$, while for $p = +\infty$ we have $\|f\|_{L^\infty} = \text{ess sup}_{x \in \mathbb{G}} |f(x)|$. Let us notice that we also have the following characterization using the distribution function $\|f\|_{L^p}^p = p \int_0^{+\infty} \alpha^{p-1} |\{x \in \mathbb{G} : |f(x)| > \alpha\}| d\alpha$.
- **weak-Lebesgue spaces** $L^{p,\infty}(\mathbb{G})$. We define them as the set of all measurable functions $f : \mathbb{G} \rightarrow \mathbb{R}$ such that $\|f\|_{L^{p,\infty}} = \sup_{\alpha > 0} \{\alpha \cdot |\{x \in \mathbb{G} : |f(x)| > \alpha\}|^{1/p}\}$ is finite. We will need the following version of Young's inequality where weak L^p spaces are involved:

Lemma 3.1 *Let $p, q, r > 1$. If $f \in L^{p,\infty}(\mathbb{G})$ and if $g \in L^r(\mathbb{G})$, then $f * g \in L^q(\mathbb{G})$ with $1 + \frac{1}{q} = \frac{1}{p} + \frac{1}{r}$ and we have the inequality $\|f * g\|_{L^q} \leq \|f\|_{L^{p,\infty}} \|g\|_{L^r}$.*

See a proof of this Lemma in [22], Theorem 1.4.24.

- **Sobolev spaces** $\dot{W}^{s,p}(\mathbb{G})$. If $1 < p < +\infty$ and for $s > 0$ we have $\|f\|_{\dot{W}^{s,p}} = \|\mathcal{J}^{\frac{s}{2}} f\|_{L^p}$, while if $p = s = 1$ we will note $\|f\|_{\dot{W}^{1,1}} = \|\nabla f\|_{L^1}$. We recall classical Sobolev inequalities in this setting:

$$\|f\|_{L^{\frac{N}{N-1}}} = \|\nabla f\|_{L^1} \quad \text{and} \quad \|f\|_{L^q} = \|f\|_{\dot{W}^{s,p}}, \quad \text{where } 1 < p < q \text{ and } -\frac{N}{q} = s - \frac{N}{p}. \quad (9)$$

- **weak Sobolev spaces** $\dot{W}_{\infty}^{s,p}(\mathbb{G})$. These spaces are defined just as classical Sobolev spaces, but we replace the L^p norm by the weak L^p one as follows:

$$\|f\|_{\dot{W}_{\infty}^{s,p}} = \|\mathcal{J}^{\frac{s}{2}} f\|_{L^{p,\infty}} \quad \text{with } 1 < p < +\infty \text{ and } s > 0.$$

- **Besov spaces** $\dot{B}_p^{s,q}(\mathbb{G})$. There are many different (and equivalent) ways to define these spaces in the setting of stratified Lie groups. In this article we will mainly use the thermic definition given by

$$\|f\|_{\dot{B}_p^{s,q}} = \left(\int_0^{+\infty} t^{(m-s/2)q} \left\| \frac{\partial^m H_t f}{\partial t^m}(\cdot) \right\|_{L^p}^q \frac{dt}{t} \right)^{1/q},$$

for $1 \leq p, q \leq +\infty, s > 0$ and m an integer such that $m > s/2$. For Besov spaces of indices $(-\beta, \infty, \infty)$ which appear in all the improved Sobolev inequalities we have:

$$\|f\|_{\dot{B}_{\infty}^{-\beta,\infty}} = \sup_{t>0} t^{\beta/2} \|H_t f\|_{L^\infty}.$$

Recall that for $0 < s < 1$ we have the inequality

$$\|\mathcal{J}^{\frac{s}{2}} f\|_{\dot{B}_{\infty}^{-\beta-s,\infty}} \leq M \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}, \quad (10)$$

where M is a universal constant.

- **Lorentz spaces** $\Lambda^p(w)(\mathbb{G})$. Let $f : \mathbb{G} \rightarrow \mathbb{R}$ be a measurable function. We define f^* , the non-increasing rearrangement of the function f , by the expression $f^*(t) = \inf\{\alpha \geq 0 : |\{x \in \mathbb{G} : |f(x)| > \alpha\}| \leq t\}$. We will say that a nonnegative locally integrable function $w : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ belongs to the Ariño-Muckenhoupt class B_p for $1 \leq p < +\infty$, if there exists $C > 0$ such that

$$\int_r^{+\infty} \left(\frac{r}{t}\right)^p w(t) dt \leq C \int_0^r w(t) dt, \quad \text{for all } 0 < r < +\infty.$$

It is not difficult to see that if $0 < p < q < +\infty$, then we have the inclusion of classes $B_p \subset B_q$. We define the Lorentz spaces $\Lambda^p(w)$ with $1 \leq p < +\infty$ by the formula

$$\|f\|_{\Lambda^p(w)} = \left(\int_0^{+\infty} (f^*(t))^p w(t) dt \right)^{\frac{1}{p}}.$$

As said in the introduction, the choice of the B_p class is due to the fact that this class of weights characterizes the boundedness of the Hardy-Littlewood maximal operator \mathcal{M}_B , given for a measurable function f by

$$\mathcal{M}_B f(x) = \sup_{B \ni x} \frac{1}{|B|} \int_B |f(y)| dy, \quad \text{where } B \text{ is an open ball,} \quad (11)$$

on the spaces $\Lambda^p(w)$ for $1 < p < +\infty$: $\|\mathcal{M}_B f\|_{\Lambda^p(w)} \leq C \|f\|_{\Lambda^p(w)}$, where C is depending on the quantity

$$[w]_{B_p} = \sup_{r>0} \left\{ r^p \left(\int_r^{+\infty} \frac{w(t)}{t^p} dt \right) / \left(\int_0^r w(t) dt \right) \right\}.$$

For more properties of these weights and the associated classical Lorentz spaces see [3], [8], [33] and [10].

- **Lorentz-Sobolev spaces** $\dot{\Lambda}^{s,p}(w)(\mathbb{G})$. Once we have fixed the base space $\Lambda^p(w)$, the homogeneous Lorentz-Sobolev spaces are easy to define and are given for $1 < p < +\infty$ and for $s > 0$ in the following way

$$\|f\|_{\dot{\Lambda}^{s,p}(w)} = \left(\int_0^{+\infty} ((\mathcal{J}^{\frac{s}{2}} f)^*(t))^p w(t) dt \right)^{\frac{1}{p}}.$$

- **weak Lorentz spaces** $\Lambda^{p,\infty}(w)(\mathbb{G})$. Let w a weight in \mathbb{R}^+ . For $0 < p < +\infty$, the weak Lorentz space $\Lambda^{p,\infty}(w)$ is the class of all measurable functions $f : \mathbb{G} \rightarrow \mathbb{R}$ such that

$$\|f\|_{\Lambda^{p,\infty}(w)} = \sup_{t>0} f^*(t) W^{1/p}(t) < +\infty,$$

where $W(t) = \int_0^t w(s) ds$. The weak Lorentz spaces were introduced in [8] and further investigated in [7], [6] and [9]. The problem of characterizing when the weak type Lorentz spaces $\Lambda^{p,\infty}(w)$, $0 < p < +\infty$ are Banach spaces was studied in [33].

- **weak Lorentz-Sobolev spaces** $\dot{\Lambda}^{s,p,\infty}(w)(\mathbb{G})$. For $1 < p < +\infty$ the homogeneous weak Lorentz-Sobolev spaces are given by

$$\|f\|_{\dot{\Lambda}^{s,p,\infty}(w)} = \sup_{t>0} (\mathcal{J}^{\frac{s}{2}} f)^*(t) \left(\int_0^t w(s) ds \right)^{1/p}.$$

- **Morrey spaces** $\mathcal{M}^{p,a}(\mathbb{G})$. For $1 < p < +\infty$ and $0 \leq a < N$, we define Morrey spaces as the space of locally integrable functions such that

$$\|f\|_{\mathcal{M}^{p,a}} = \sup_{x_0 \in \mathbb{R}^n} \sup_{0 < r < +\infty} \left(\frac{1}{r^a} \int_{B(x_0,r)} |f(x)|^p dx \right)^{\frac{1}{p}} < +\infty.$$

Morrey spaces are indeed a generalization of Lebesgue spaces since when $a = 0$ we have $\mathcal{M}^{p,0} \simeq L^p$. The use of Morrey and Morrey-Sobolev spaces in this article is due to the fact that the Hardy-Littlewood maximal operator (11) is also bounded in such spaces. See more details in [14] in the framework of \mathbb{R}^n and [2] in the setting of stratified Lie groups. See also [32] and the references given there in for other interesting generalizations.

- **Morrey-Sobolev spaces** $\dot{\mathcal{M}}^{s,p,a}(\mathbb{G})$. For $0 < s$ and $1 < p < +\infty$ with $0 \leq a < N$ we consider the homogeneous Morrey-Sobolev spaces $\dot{\mathcal{M}}^{s,p,a}$ by the quantity

$$\|f\|_{\dot{\mathcal{M}}^{s,p,a}} = \|\mathcal{J}^{\frac{s}{2}}f\|_{\mathcal{M}^{p,a}}.$$

4 Proof of Theorem 1

We will prove here, in the framework of stratified Lie groups, the inequality

$$\|f\|_{\dot{\Lambda}^{s_1,q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s,p}(w)}^\theta \|f\|_{\dot{B}_\infty^{-\beta,\infty}}^{1-\theta},$$

where $f : \mathbb{G} \rightarrow \mathbb{R}$ is a function such that $f \in \dot{\Lambda}^{s,p}(w)(\mathbb{G}) \cap \dot{B}_\infty^{-\beta,\infty}(\mathbb{G})$ with $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$. We will always assume here that w is a weight in the Ariño-Muckenhoupt class B_p . The reason for this particular choice of weights relies on the fact that we will need the boundedness of the Hardy-Littlewood maximal operator on Lorentz $\Lambda^p(w)$ spaces and this is ensured by the condition $w \in B_p$. See [3] and [10] for details.

By the definition of Lorentz-Sobolev spaces given in Section 3, this inequality can be rewritten in the following way

$$\|\mathcal{J}^{\frac{s_1}{2}}f\|_{\Lambda^q(w)} \leq C \|\mathcal{J}^{\frac{s}{2}}f\|_{\Lambda^p(w)}^\theta \|f\|_{\dot{B}_\infty^{-\beta,\infty}}^{1-\theta}.$$

For the proof of this inequality, we will use a variant of Hedberg's inequality. Indeed, since $0 < s_1 < s$, we use the characterization of the positive powers of the Laplacian given in (8) and we have for $k > s/2 > s_1/2$

$$\begin{aligned} \mathcal{J}^{\frac{s_1}{2}}f(x) &= \frac{1}{\Gamma(k - s_1/2)} \int_0^{+\infty} t^{k - \frac{s_1}{2} - 1} \mathcal{J}^k H_t f(x) dt \\ &= \frac{1}{\Gamma(k - s_1/2)} \left(\int_0^T t^{k - \frac{s_1}{2} - 1} \mathcal{J}^k H_t f(x) dt + \int_T^{+\infty} t^{k - \frac{s_1}{2} - 1} \mathcal{J}^k H_t f(x) dt \right), \end{aligned}$$

where T will be defined below. In particular we have

$$|\mathcal{J}^{\frac{s_1}{2}}f(x)| \leq \frac{1}{\Gamma(k - s_1/2)} \left(\int_0^T t^{k - \frac{s_1}{2} - 1} |\mathcal{J}^k H_t f(x)| dt + \int_T^{+\infty} t^{k - \frac{s_1}{2} - 1} |\mathcal{J}^k H_t f(x)| dt \right). \quad (12)$$

For the first integral of the right-hand side of the previous formula we will use the following fact.

Lemma 4.1 *Let $f \in \mathcal{S}'(\mathbb{G})$ and $\varphi \in \mathcal{S}(\mathbb{G})$. We denote by $\mathcal{M}_\varphi(f)$ the maximal function of f (with respect to φ) which is given by the expression*

$$\mathcal{M}_\varphi f(x) = \sup_{0 < t < +\infty} \{|f * \varphi_t(x)|\}, \quad \text{with } \varphi_t(x) = t^{-N/2} \varphi(t^{-1/2}x).$$

If the function φ is such that $|\varphi(x)| \leq C(1 + |x|)^{-N-\varepsilon}$ for some $\varepsilon > 0$, then we have the following pointwise inequality

$$\mathcal{M}_\varphi f(x) \leq C\mathcal{M}_B f(x),$$

where $\mathcal{M}_B f(x)$ is the Hardy-Littlewood maximal function defined by (11).

For a proof of this lemma see [22] or [18]. With this lemma in mind, and since $k > s/2$, we remark that we have the identity $\mathcal{J}^k H_t f(x) = \mathcal{J}^{k-\frac{s}{2}} h_t * \mathcal{J}^{\frac{s}{2}} f(x)$. Now, by homogeneity we obtain $\mathcal{J}^{k-\frac{s}{2}}(h_t)(x) = t^{-k+\frac{s}{2}}(\mathcal{J}^{k-\frac{s}{2}} h_t)(x)$ and if we denote φ_t by $\varphi_t(x) = (\mathcal{J}^{k-\frac{s}{2}} h_t)(x)$ we have that $\varphi_t(x) = t^{-N/2}\varphi(t^{-1/2}x)$, moreover, since the heat kernel h_t is a smooth function, with the previous notation we obtain $|\varphi(x)| \leq C(1 + |x|)^{-N-\varepsilon}$. Then we can write

$$\mathcal{J}^k H_t f(x) = t^{-k+\frac{s}{2}} \varphi_t * \mathcal{J}^{\frac{s}{2}} f(x),$$

and applying the Lemma 4.1 we have the following pointwise inequality for the first term of (12):

$$|\mathcal{J}^k H_t f(x)| = t^{-k+\frac{s}{2}} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right) (x).$$

Now, for the second integral of the right-hand side of (12) we simply use the fact that $\|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}} \simeq \|f\|_{\dot{B}_\infty^{-\beta,\infty}}$ and the thermic definition of Besov spaces to obtain

$$|\mathcal{J}^k H_t f(x)| = |H_t \mathcal{J}^k f(x)| \leq C t^{\frac{-\beta-2k}{2}} \|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}}.$$

With these two inequalities at hand, we apply them in (12) and one has

$$\begin{aligned} |\mathcal{J}^{\frac{s_1}{2}} f(x)| &\leq \frac{C}{\Gamma(k - s_1/2)} \left(\int_0^T t^{k-\frac{s_1}{2}-1} t^{-k+\frac{s}{2}} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right) (x) dt + \int_T^{+\infty} t^{k-\frac{s_1}{2}-1} t^{\frac{-\beta-2k}{2}} \|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}} dt \right) \\ &\leq \frac{C}{\Gamma(k - s_1/2)} \left(T^{\frac{s-s_1}{2}} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right) (x) + T^{\frac{-\beta-s_1}{2}} \|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}} \right). \end{aligned}$$

We fix now the parameter T by the condition

$$T = \left(\frac{\|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}}}{\mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right) (x)} \right)^{\frac{2}{\beta+s}},$$

and we obtain the following inequality

$$|\mathcal{J}^{\frac{s_1}{2}} f(x)| \leq \frac{C}{\Gamma(k - s_1/2)} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right)^{1-\frac{s-s_1}{\beta+s}} (x) \|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}}^{\frac{s-s_1}{\beta+s}}.$$

Since $\frac{s-s_1}{\beta+s} = 1 - \theta$ and using again the fact $\|\mathcal{J}^k f\|_{\dot{B}_\infty^{-\beta-2k,\infty}} \simeq \|f\|_{\dot{B}_\infty^{-\beta,\infty}}$ we have

$$|\mathcal{J}^{\frac{s_1}{2}} f(x)| \leq \frac{C}{\Gamma(k - s_1/2)} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right)^\theta (x) \|f\|_{\dot{B}_\infty^{-\beta,\infty}}^{1-\theta}. \quad (13)$$

Once we have obtained this pointwise inequality, we will use the following properties of the non-increasing rearrangement function.

Lemma 4.2 *If $f, g : \mathbb{G} \rightarrow \mathbb{R}$ are two measurable functions, we have*

- 1) *if $|g| \leq |f|$ a.e. then $g^* \leq f^*$,*
- 2) *if $0 < \theta$, then $(|f|^\theta)^* = (f^*)^\theta$.*

For a proof see Proposition 1.4.5 of [22]. Recalling that $\theta = p/q$ and applying these facts to the inequality (13) we obtain

$$\left((\mathcal{J}^{\frac{s_1}{2}} f)^*(t) \right)^q \leq C \left((\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f))^*(t) \right)^p \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p}. \quad (14)$$

Multiplying the previous inequality by a weight w from the Ariño-Muckenhoupt class B_p and integrating with respect to the variable t we obtain

$$\int_0^{+\infty} \left((\mathcal{J}^{\frac{s_1}{2}} f)^*(t) \right)^q w(t) dt \leq C \int_0^{+\infty} \left((\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f))^*(t) \right)^p w(t) dt \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p},$$

and then, by the definition of classical Lorentz spaces given in Section 3 we have

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^q(w)} \leq C \|\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f)\|_{\Lambda^p(w)}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}.$$

Now, since the weight w belongs to the class B_p with $1 < p < +\infty$, we have that the Hardy-Littlewood maximal operator is bounded on the space $\Lambda^p(w)$ and we obtain

$$\|\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f)\|_{\Lambda^p(w)} \leq \|\mathcal{J}^{\frac{s}{2}} f\|_{\Lambda^p(w)},$$

and finally we have the desired inequality for classical Lorentz spaces:

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^q(w)} \leq C \|\mathcal{J}^{\frac{s}{2}} f\|_{\Lambda^p(w)}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}.$$

■

Now we will state in the following corollaries some interesting consequences of this previous theorem.

Corollary 4.1 *Let $w \in B_p$ be a weight and let $f : \mathbb{G} \rightarrow \mathbb{R}$ be a function such that $f \in \dot{\Lambda}^{s,p,\infty}(w)(\mathbb{G}) \cap \dot{B}_{\infty}^{-\beta,\infty}(\mathbb{G})$. Then we have the following version of improved Sobolev inequalities of weak type:*

$$\|f\|_{\dot{\Lambda}^{s_1,q,\infty}(w)} \leq C \|f\|_{\dot{\Lambda}^{s,p,\infty}(w)}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta},$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

Proof. We start again with the pointwise inequality (14):

$$\left((\mathcal{J}^{\frac{s_1}{2}} f)^*(t) \right)^q \leq C \left((\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f))^*(t) \right)^p \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p}.$$

Now, we multiply both parts of this inequality by $W(t)$ and we take the supremum in the variable t :

$$\begin{aligned} \|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^q(w)}^q &= \sup_{t>0} W(t) \left((\mathcal{J}^{\frac{s_1}{2}} f)^*(t) \right)^q \leq C \sup_{t>0} \left\{ \left((\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f))^*(t) \right)^p W(t) \right\} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p} \\ &\leq C \|\mathcal{M}_B (\mathcal{J}^{\frac{s}{2}} f)\|_{\Lambda^p(w)}^p \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p}, \end{aligned}$$

since it is known (see e.g. [33]) that for $w \in B_p$ the Hardy-Littlewood maximal operator \mathcal{M}_B is bounded on $\Lambda^{p,\infty}(w)$, therefore we obtain that

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^{q,\infty}(w)} \leq C \|\mathcal{J}^{\frac{s}{2}} f\|_{\Lambda^{p,\infty}(w)}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta}.$$

■

Now we will study other variations of the previous results by considering a different type of weights. To be more precise, we will study two-weighted inequalities and in what follows, for v and w two weights and for $t > 0$, we will denote by $V(t)$ and $W(t)$ the quantities $V(t) = \int_0^t v(s)ds$ and $W(t) = \int_0^t w(s)ds$.

Our first two-weighted improved Lorentz-Sobolev inequality is given in the following corollary.

Corollary 4.2 *Let $1 < p < q < +\infty$ and let (v, w) be a pair of positive weights satisfying the following properties*

$$\sup_{t>0} \frac{W(t)^{1/p}}{V(t)^{1/p}} < +\infty \quad \text{and} \quad \sup_{t>0} \left(\int_t^{+\infty} \frac{w(s)}{s^p} ds \right)^{1/p} \left(\int_0^t \frac{v(s)s^{p'}}{V(s)^{p'}} ds \right)^{1/p'} < +\infty.$$

If $f : \mathbb{G} \rightarrow \mathbb{R}$ is a function such that $f \in \dot{\Lambda}^{s,p}(v) \cap \dot{B}_{\infty}^{-\beta,\infty}$ with $s > 0$, then we have a two-weighted version of improved Sobolev inequalities

$$\|f\|_{\dot{\Lambda}^{s_1,q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s,p}(v)}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta},$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

This inequality is interesting since it is possible, under some hypotheses, to consider different weights in the left-hand side and in the right-hand side of the inequality.

Proof. Using the pointwise inequality (14) and the fact that the Hardy-Littlewood maximal operator

$$\mathcal{M}_B : \Lambda^p(v) \rightarrow \Lambda^p(w)$$

is bounded for such weights (see [37] for details) we obtain the desired inequality. ■

If we are allowed to change the weights that define the Lorentz spaces in the previous inequalities, it is then also possible to change, with specific conditions on the weights, the parameters of these spaces. In the following corollary we gather some results where we consider different Lorentz spaces in the right-hand side of the inequality. Indeed, the first point is a generalization of the previous corollary and we will consider in the right-hand side Lorentz-Sobolev spaces of type $\dot{\Lambda}^{s,q_0}(v)$ instead of $\dot{\Lambda}^{s,p}(v)$ where $1 < q_0 \leq p < +\infty$. The second point allows us to study the case when $1 < p < q_0 < +\infty$ and finally, the third point treats the case when $0 < q_0 < 1$.

Corollary 4.3 *Let $0 < q_0 < +\infty$, $s > 0$, let $f : \mathbb{G} \rightarrow \mathbb{R}$ be a measurable function and let (v, w) be a pair of weights.*

1) If $1 < q_0 \leq p < +\infty$ and if (v, w) are satisfying the following conditions

$$\sup_{t>0} \frac{W(t)^{1/p}}{V(t)^{1/q_0}} < +\infty \quad (15)$$

and

$$\sup_{t>0} \left(\int_0^t \frac{w(s)}{s^p} ds \right)^{1/p} \left(\int_0^t \frac{v(s)s^{q_0'}}{V(s)^{q_0}} \right) < +\infty, \quad (16)$$

then, if $f \in \dot{\Lambda}^{s, q_0}(v)(\mathbb{G}) \cap \dot{B}_{\infty}^{-\beta, \infty}(\mathbb{G})$, we have the following inequality

$$\|f\|_{\dot{\Lambda}^{s_1, q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s, q_0}(v)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta},$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

2) If $1 < p < q_0 < +\infty$ and (v, w) are satisfying

$$\left(\int_0^{+\infty} \left(\frac{W(s)}{V(s)} \right)^{r/q_0} w(s) ds \right)^{1/r} < +\infty$$

and

$$\left(\int_0^{+\infty} \left[\left(\int_s^{+\infty} \frac{w(t)}{t^p} dt \right)^{1/p} \left(\int_0^t \frac{v(t)t^{q_0'}}{V(t)^{q_0'}} dt \right)^{1/p'} \right]^r \frac{v(s)s^{q_0'}}{V(s)^{q_0'}} ds \right)^{1/r} < +\infty,$$

where r is given by $\frac{1}{r} = \frac{1}{p} - \frac{1}{q}$ and $\frac{1}{q_0} + \frac{1}{q_0'} = 1$. Then, if $f \in \dot{\Lambda}^{s, q_0}(v)(\mathbb{G}) \cap \dot{B}_{\infty}^{-\beta, \infty}(\mathbb{G})$, we have

$$\|f\|_{\dot{\Lambda}^{s_1, q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s, q_0}(v)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta},$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

3) If $0 < q_0 < 1$ and $1 < p < +\infty$ and if (v, w) are satisfying (15) and

$$\sup_{t>0} \frac{t}{V(t)^{1/q_0}} \left(\int_t^{+\infty} \frac{w(s)}{s^p} ds \right)^{1/p} < +\infty,$$

then, assuming that $f \in \dot{\Lambda}^{s, q_0}(v)(\mathbb{G}) \cap \dot{B}_{\infty}^{-\beta, \infty}(\mathbb{G})$, we obtain

$$\|f\|_{\dot{\Lambda}^{s_1, q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s, q_0}(v)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta},$$

where $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

Proof. From the pointwise inequality (14) we obtain that

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^q(w)} \leq C \|\mathcal{M}_B(\mathcal{J}^{\frac{s}{2}} f)\|_{\Lambda^p(w)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}.$$

Now, under all these hypotheses on the weights v and w , we have that the Hardy-Littlewood maximal operator $\mathcal{M}_B : \Lambda^{q_0}(v) \rightarrow \Lambda^p(w)$ is bounded (see [37] and [7]) and then we obtain

$$\|f\|_{\dot{\Lambda}^{s_1, q}(w)} \leq C \|f\|_{\dot{\Lambda}^{s, q_0}(v)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}.$$

■

We have also the following two-weighted version of improved Sobolev inequalities of weak type:

Corollary 4.4 *Let $1 < p < +\infty$, $0 < q_0 < +\infty$. Let (v, w) be a pair of weights such that*

$$\sup_{t>0} \frac{W(t)^{1/p}}{t} \int_0^t V^{-1/q_0}(s) ds < +\infty, \quad (17)$$

and let $f : \mathbb{G} \rightarrow \mathbb{R}$ be a function such that $f \in \dot{\Lambda}^{s, q_0, \infty}(v)(\mathbb{G}) \cap \dot{B}_{\infty}^{-\beta, \infty}(\mathbb{G})$. Then we have the following inequality

$$\|f\|_{\dot{\Lambda}^{s_1, q, \infty}(w)} \leq C \|f\|_{\dot{\Lambda}^{s, q_0, \infty}(v)}^{\theta} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta},$$

where $0 < q_0 < +\infty$, $1 < p < q < +\infty$, $\theta = p/q$, $s_1 = \theta s - (1 - \theta)\beta$ and $-\beta < s_1 < s$.

Proof. It is enough to follow the same lines of the Corollary 4.1 to obtain

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^{q, \infty}(w)}^q \leq C \|\mathcal{M}_B(\mathcal{J}^{\frac{s}{2}} f)\|_{\Lambda^{p, \infty}(w)}^p \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p},$$

since the pair of weights (v, w) satisfies the condition (17) it implies that the operator $\mathcal{M}_B : \Lambda^{q_0, \infty}(v) \rightarrow \Lambda^{p, \infty}(w)$ is bounded (see [33]) and we obtain

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\Lambda^{q, \infty}(w)}^q \leq C \|\mathcal{J}^{\frac{s}{2}} f\|_{\Lambda^{q, \infty}(v)}^p \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q-p},$$

which is the desired inequality. ■

5 Generalizations

In this section we give some generalizations of Theorems 1 and we prove Theorem 2. These generalizations are made possible since the techniques developed in our proofs are based on general harmonic analysis arguments and since many of the tools used in this article are available in other frameworks. Indeed, the spectral theory associated to the Laplace operator, the boundedness of the Hardy-Littlewood maximal operator and the use of appropriate weights in order to define well suited functional spaces are intensively studied and many interesting properties were generalized to different settings.

5.1 Morrey spaces

We prove now Theorem 2 in the setting of stratified Lie groups. Morrey spaces were studied in this framework by many authors, see for example the articles [2], [30] and the references there in.

As said in the introduction, once we have at our disposal the fact that the Hardy-Littlewood maximal operator is bounded in the convenient functional framework, it is possible to improve Sobolev inequalities in the following way. The starting point of our proof is the pointwise inequality (13):

$$|\mathcal{J}^{\frac{s_1}{2}} f(x)| \leq \frac{C}{\Gamma(k - s_1/2)} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right)^{\theta}(x) \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}.$$

Since $\theta = p/q$ we have for $r > 0$ and for $0 \leq a < N$ the inequalities

$$\begin{aligned} \frac{1}{r^a} \int_{B(x_0, r)} |\mathcal{J}^{\frac{s_1}{2}} f(x)|^q dx &\leq C \left(\frac{1}{r^a} \int_{B(x_0, r)} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right)^p(x) dx \right) \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{q(1-\theta)} \\ \left(\frac{1}{r^a} \int_{B(x_0, r)} |\mathcal{J}^{\frac{s_1}{2}} f(x)|^q dx \right)^{1/q} &\leq C \left(\frac{1}{r^a} \int_{B(x_0, r)} \mathcal{M}_B \left(\mathcal{J}^{\frac{s}{2}} f \right)^p(x) dx \right)^{1/q} \|f\|_{\dot{B}_{\infty}^{-\beta, \infty}}^{1-\theta}, \end{aligned}$$

from which we derive the estimate

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\mathcal{M}^{q,a}} \leq C \|\mathcal{M}_B(\mathcal{J}^{\frac{s}{2}} f)\|_{\mathcal{M}^{p,a}}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta}.$$

In order to conclude, we use the fact that the Hardy-Littlewood maximal operator is bounded in Morrey spaces and we obtain

$$\|\mathcal{J}^{\frac{s_1}{2}} f\|_{\mathcal{M}^{q,a}} \leq C \|\mathcal{J}^{\frac{s}{2}} f\|_{\mathcal{M}^{p,a}}^\theta \|f\|_{\dot{B}_{\infty}^{-\beta,\infty}}^{1-\theta},$$

which is the desired inequality stated in Theorem 2.

Remark 5.1 *The boundedness of the Hardy-Littlewood maximal operator was studied for generalized Morrey spaces in [2], [29] and [32]. As long as this boundedness property is satisfied it should be possible to generalize Theorem 2. Indeed, from the pointwise inequality (13) it should be easy (taking into account the necessary precautions) to reconstruct the corresponding norms in order to obtain an improved Sobolev-like inequality.*

5.2 Nilpotent Lie groups

We consider now a more general framework than the one given by stratified Lie groups. Indeed, going one step further in the process of generalization, it is possible to consider nilpotent Lie groups since all the tools used in the proof of Theorem 1 are available in these settings.

We recall for the sake of completeness this framework. Let \mathbb{G} be a connected unimodular Lie group endowed with its Haar measure dx . Denote by \mathfrak{g} the Lie algebra of \mathbb{G} and consider a family (that will be fixed from now on) of left-invariant vector fields on \mathbb{G}

$$\mathbf{X} = \{X_1, \dots, X_k\},$$

satisfying the *Hörmander condition*². We endow the group \mathbb{G} with a metric structure by considering the Carnot-Carathéodory metric associated with \mathbf{X} . See [38] for details. We will denote $\|x\|$ the distance between the origin e and x and $\|y^{-1} \cdot x\|$ the distance between x and y . For $r > 0$ and $x \in \mathbb{G}$, denote by $B(x, r)$ the open ball with respect to the Carnot-Carathéodory metric centered in x and of radius r , and by $V(r) = \int_{B(x,r)} dx$ the Haar measure of any ball of radius r . When $0 < r < 1$, there exists $d \in \mathbb{N}^*$, c_l and $C_l > 0$ such that, for all $0 < r < 1$ we have

$$c_l r^d \leq V(r) \leq C_l r^d.$$

The integer d is the *local dimension* of (\mathbb{G}, \mathbf{X}) . When $r \geq 1$, two situations may occur, independently of the choice of the family \mathbf{X} : either \mathbb{G} has polynomial volume growth and there exist $D \in \mathbb{N}^*$, c_∞ and $C_\infty > 0$ such that, for all $r \geq 1$ we have

$$c_\infty r^D \leq V(r) \leq C_\infty r^D,$$

or \mathbb{G} has exponential volume growth, which means that there exist $c_e, C_e, \alpha, \beta > 0$ such that, for all $r \geq 1$ we have

$$c_e e^{\alpha r} \leq V(r) \leq C_e e^{\beta r}.$$

²which means that the Lie algebra generated by the family \mathbf{X} is \mathfrak{g} .

When \mathbb{G} has polynomial volume growth, the integer D is called the *dimension at infinity* of \mathbb{G} . Recall that nilpotent groups have polynomial volume growth and that a *strict* subclass of the nilpotent groups consists of stratified Lie groups where $d = D$.

Once we have fixed the family \mathbf{X} , we define the gradient on \mathbb{G} by $\nabla = (X_1, \dots, X_k)$ and we consider a Laplacian \mathcal{J} on \mathbb{G} defined in the same way as in (7)

$$\mathcal{J} = - \sum_{j=1}^k X_j^2,$$

which is a positive self-adjoint, hypo-elliptic operator since \mathbf{X} satisfies the Hörmander's condition, see [38]. Its associated heat operator on $]0, +\infty[\times \mathbb{G}$ is given by $\partial_t + \mathcal{J}$ and we will denote by $(H_t)_{t>0}$ the semi-group obtained from the Laplacian \mathcal{J} . It is worth noting that many of the properties given in Theorem 3 remain true for the heat semi-group H_t in this general setting. For more details concerning nilpotent Lie groups see the books [38], [18], [35] and the articles [19], [31], [12] and the references there in. Fractional powers of the Laplacian can be defined in a completely similar way using the expression (8). It is then possible to define all the functional spaces given in Section 3 in the framework of nilpotent Lie groups.

With all these preliminaries, we see that we have at our disposal all the ingredients needed in order to perform the computations done in Sections 4, and thus Theorem 1 can be generalized to the setting of nilpotent Lie groups.

Acknowledgments. A part of this work was performed while the second and third authors visited the University of Evry Val d'Essonne. We express our gratitude to the Laboratoire de Mathématiques et Modélisation d'Evry (LaMME) of the University of Evry Val d'Essonne for the hospitality and excellent conditions. The second named author was partially supported by POSDRU/159/1.5/S/137750.

References

- [1] D. ADAMS. *A note on Riesz potentials*. Duke Math. J., 42 (1975), no. 4, 765–778.
- [2] A. AKBULUT, V. GULIYEV, Y. MAMMADOV. *Boundedness of fractional maximal operator and their higher order commutator in the generalized Morrey spaces on Carnot groups*. Acta Math. Sci. Ser. B Engl. Ed. 33 (2013), no. 5, 1329–1346.
- [3] M. ARIÑO, B. MUCKENHOUPT. *Maximal functions on classical Lorentz spaces and Hardy's inequality with weights for nonincreasing functions*. Trans. Amer. Math. Soc. 320 (1990), no. 2, 727–735.
- [4] H. BAHOURI, A. COHEN. *Refined Sobolev inequalities in Lorentz spaces*. J. Fourier Anal. Appl. 17 (2011), no. 4, 662–673.
- [5] C. BENNETT, R. SHARPLEY. *Interpolation of operators*. Pure and Applied Mathematics, Vol 129, Academic Press Inc., Boston, MA, (1988).
- [6] M. J. CARRO, A. GARCÍA DEL AMO, J. SORIA. *Weak-type weights and normable Lorentz spaces*. Proc. Amer. Math. Soc. 124 (1996), no. 3, 849–857.
- [7] M. J. CARRO, J. SORIA. *Boundedness of some integral operators*. Canad. J. Math. 45 (1993), no. 6, 1155–1166.

- [8] M. J. CARRO, J. SORIA. *Weighted Lorentz spaces and the Hardy operator*. J. Funct. Anal. 112 (1993), no. 2, 480–494.
- [9] M. J. CARRO, J. SORIA. *The Hardy-Littlewood maximal function and weighted Lorentz spaces*. J. London Math. Soc. (2) 55 (1997), no. 1, 146–158.
- [10] M. J. CARRO, J.A. RAPOSO, J. SORIA. *Recent developments in the theory of Lorentz spaces and weighted inequalities*. Mem. Amer. Math. Soc. 187 (2007), no. 877.
- [11] D. CHAMORRO, *Improved Sobolev Inequalities and Muckenhoupt weights on stratified Lie groups*. J. Math. Anal. Appl. 377 (2011), no. 2, 695–709.
- [12] D. CHAMORRO, *Some functional inequalities on polynomial volume growth Lie groups*, Canad. J. Math. 64 (2012), 481–496.
- [13] D. CHAMORRO, P.G. LEMARIÉ-RIEUSSET. *Real interpolation method, Lorentz spaces and refined Sobolev inequalities*. J. Funct. Anal. 265 (2013), no. 12, 3219–3232.
- [14] F. CHIARENZA, M. FRASCA. *Morrey spaces and Hardy-Littlewood maximal function*. Rend. Mat. Appl. (7) 7 (1987), no. 3-4, 273–279.
- [15] A. CIANCHI. *Optimal Orlicz-Sobolev embeddings*. Rev. Mat. Iberoamericana 20 (2004), no. 2, 427–474.
- [16] A. COHEN, W. DAHMEN, I. DAUBECHIES & R. DE VORE. *Harmonic Analysis of the space BV*. Rev. Mat. Iberoamericana 19 (2003), no. 1, 235–263.
- [17] G. FOLLAND. *Subelliptic estimates and function spaces on nilpotent Lie groups*. Ark. Mat. 13 (1975), no. 2, 161–207.
- [18] G. FOLLAND & E. M. STEIN. *Hardy Spaces on homogeneous groups*. Mathematical Notes, 28, Princeton University Press (1982).
- [19] G. FURIOLI, C. MELZI & A. VENERUSO. *Littlewood-Paley decomposition and Besov spaces on Lie groups of polynomial growth*. Math. Nachr. 279 (2006), no. 9-10, 1028–1040.
- [20] N. GAROFALO & D NHIEU. *Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces*. Comm. Pure Appl. Math. 49 (1996), no. 10, 1081–1144.
- [21] P. GÉRARD, Y. MEYER & F. ORU. *Inégalités de Sobolev Précisées*. Séminaire sur les Equations aux Dérivées Partielles, 1996-1997, Exp. No. IV, École Polytech., Palaiseau.
- [22] L. GRAFAKOS. *Classical and Modern Fourier Analysis*. Prentice Hall (2004).
- [23] V. I. KOLYADA, F. J. PÉREZ LÁZARO. *On Gagliardo-Nirenberg Type Inequalities*. J. Fourier Anal. Appl. 20 (2014), No. 3, 577–607.
- [24] M. LEDOUX. *On improved Sobolev embedding theorems*. Math. Res. Lett. 10 (2003), no. 5-6, 659–669.
- [25] G. G. LORENTZ. *Some new functional spaces*, Ann. of Math. (2) 51 (1950), 37–55.
- [26] G. G. LORENTZ. *On the theory of spaces Λ* , Pacific J. Math. 1 (1951), 411–429.
- [27] J. MARTN, M. MILMAN, *Sharp Gagliardo-Nirenberg inequalities via symmetrization*. Math. Res. Lett. 14 (2007), no. 1, 49–62.
- [28] B. MUCKENHOUPT. *Weighted norm inequalities for the Hardy maximal function*. Trans. Amer. Math. Soc. 165 (1972), 207–226.
- [29] E. NAKAI, *Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*. Math. Nachr. 166 (1994), 95–103.

- [30] E. NAKAI, *The Campanato, Morrey and Hölder spaces on spaces of homogeneous type*. Studia Math. 176 (2006), no.1, 1–19.
- [31] K. SAKA. *Besov spaces and Sobolev spaces on a nilpotent Lie group*. Tôhoku. Math. J. (2) 31 (1979), no. 4, 383–437.
- [32] Y. SAWANO, S. SUGANO, H. TANAKA, *Generalized fractional integral operators and fractional maximal operators in the framework of Morrey spaces*, Trans. Amer. Math. Soc. 363 (2011), no. 12, 6481–6503.
- [33] J. SORIA. *Lorentz spaces of weak-type*. Quart. J. Math. Oxford Ser. (2) 49 (1998), no. 193, 93–103.
- [34] E. M. STEIN. *Singular Integrals and Differentiability Properties of Functions*. Princeton Mathematical Series, 30. Princeton University Press (1970).
- [35] E. M. STEIN. *Harmonic Analysis*. Princeton University Press (1993).
- [36] R. STRICHARTZ. *Self-similarity on nilpotent Lie groups*. Contemp. Math., 140 (1992), 123–157.
- [37] E. SAWYER. *Boundedness of classical operators on classical Lorentz spaces*. Studia Math. 96 (1990), no. 2, 145–158.
- [38] N. Th. VAROPOULOS, L. SALOFF-COSTE & T. COULHON. *Analysis and geometry on groups*. Cambridge Tracts in Mathematics, 100 (1992).

Diego CHAMORRO

Laboratoire de Mathématiques et Modélisation d'Evry
(LaMME) UMR 8071, Université d'Evry Val d'Essonne
23 Boulevard de France, 91037 Evry Cedex
France
diego.chamorro@univ-evry.fr

Anca MARCOCI & Liviu MARCOCI

Department of Mathematics and Computer Science
Technical University of Civil Engineering
Bucharest, Bld. Lacul Tei, no. 124, sector 2.
Romania
anca.marcoci@utcb.ro lmarcoci@instal.utcb.ro